

Exercise 13

In Exercises 11-14, (a) solve the given equation by the method of characteristic curves, and (b) check your answer by plugging it back into the equation.

$$\frac{\partial u}{\partial x} + \sin x \frac{\partial u}{\partial y} = 0.$$

Solution

The differential of a two-dimensional function $g = g(x, y)$ is given by

$$dg = \frac{\partial g}{\partial x} dx + \frac{\partial g}{\partial y} dy.$$

Dividing both sides by dx yields the fundamental relationship between the total derivative of g and its partial derivatives.

$$\frac{dg}{dx} = \frac{\partial g}{\partial x} + \frac{dy}{dx} \frac{\partial g}{\partial y}$$

Comparing this to the PDE, we see that along the (characteristic) curves in the xy -plane defined by

$$\frac{dy}{dx} = \sin x \tag{1}$$

the PDE reduces to the ODE,

$$\frac{du}{dx} = 0. \tag{2}$$

Solve equation (1), using ξ for the characteristic coordinate.

$$y = -\cos x + \xi \quad \rightarrow \quad \xi = y + \cos x$$

Then solve equation (2) by integrating both sides with respect to x .

$$u(x, \xi) = f(\xi)$$

Here f is an arbitrary function. Now that u is known, change back to the original variables.

$$u(x, y) = f(y + \cos x)$$

Compute the first derivatives to check the solution.

$$\frac{\partial u}{\partial x} = f'(y + \cos x) \cdot \frac{\partial}{\partial x}(y + \cos x) = f'(y + \cos x) \cdot (-\sin x) = -(\sin x)f'$$

$$\frac{\partial u}{\partial y} = f'(y + \cos x) \cdot \frac{\partial}{\partial y}(y + \cos x) = f'(y + \cos x) \cdot (1) = f'$$

As a result,

$$\frac{\partial u}{\partial x} + \sin x \frac{\partial u}{\partial y} = -(\sin x)f' + (\sin x)f' = 0.$$