

## Exercise 8

Prove that two nonzero complex numbers  $z_1$  and  $z_2$  have the same moduli if and only if there are complex numbers  $c_1$  and  $c_2$  such that  $z_1 = c_1 c_2$  and  $z_2 = c_1 \bar{c}_2$ .

*Suggestion:* Note that

$$\exp\left(i\frac{\theta_1 + \theta_2}{2}\right) \exp\left(i\frac{\theta_1 - \theta_2}{2}\right) = \exp(i\theta_1)$$

and

$$\exp\left(i\frac{\theta_1 + \theta_2}{2}\right) \overline{\exp\left(i\frac{\theta_1 - \theta_2}{2}\right)} = \exp(i\theta_2).$$

### Solution

Suppose that there are two nonzero complex numbers,  $z_1$  and  $z_2$ ,

$$z_1 = r_1 e^{i\theta_1} \quad z_2 = r_2 e^{i\theta_2}$$

and that they have the same moduli.

$$|z_1| = |z_2| \quad \rightarrow \quad |r_1 e^{i\theta_1}| = |r_2 e^{i\theta_2}| \quad \rightarrow \quad r_1 = r_2$$

If we choose  $c_1$  and  $c_2$  to be

$$c_1 = r_1 \exp\left(i\frac{\theta_1 + \theta_2}{2}\right)$$

$$c_2 = \exp\left(i\frac{\theta_1 - \theta_2}{2}\right),$$

then

$$z_1 = c_1 c_2 = r_1 \exp\left(i\frac{\theta_1 + \theta_2}{2}\right) \exp\left(i\frac{\theta_1 - \theta_2}{2}\right) = r_1 e^{i\theta_1}$$

$$z_2 = c_1 \bar{c}_2 = r_1 \exp\left(i\frac{\theta_1 + \theta_2}{2}\right) \overline{\exp\left(i\frac{\theta_1 - \theta_2}{2}\right)} = r_1 e^{i\theta_2} = r_2 e^{i\theta_2}.$$

The first part of the proof is complete. Suppose now that there exist complex numbers,  $c_1$  and  $c_2$ , such that  $z_1 = c_1 c_2$  and  $z_2 = c_1 \bar{c}_2$ . The aim is to show that  $z_1$  and  $z_2$  have the same moduli. If we have

$$c_1 = a_1 + ib_1$$

$$c_2 = a_2 + ib_2,$$

then

$$z_1 = c_1 c_2 = (a_1 + ib_1)(a_2 + ib_2) = a_1 a_2 - b_1 b_2 + i(a_1 b_2 + a_2 b_1)$$

$$z_2 = c_1 \bar{c}_2 = (a_1 + ib_1)(a_2 - ib_2) = a_1 a_2 + b_1 b_2 + i(-a_1 b_2 + a_2 b_1),$$

and the magnitudes are

$$|z_1| = \sqrt{(a_1 a_2 - b_1 b_2)^2 + (a_1 b_2 + a_2 b_1)^2} = \sqrt{a_1^2 a_2^2 + a_2^2 b_1^2 + a_1^2 b_2^2 + b_1^2 b_2^2}$$

$$|z_2| = \sqrt{(a_1 a_2 + b_1 b_2)^2 + (-a_1 b_2 + a_2 b_1)^2} = \sqrt{a_1^2 a_2^2 + a_2^2 b_1^2 + a_1^2 b_2^2 + b_1^2 b_2^2}$$

in fact equal. Therefore, two nonzero complex numbers,  $z_1$  and  $z_2$ , have the same moduli if and only if there are complex numbers,  $c_1$  and  $c_2$ , such that  $z_1 = c_1 c_2$  and  $z_2 = c_1 \bar{c}_2$ .