

## Exercise 16

Let  $L_n$  denote the left-endpoint sum using  $n$  subintervals and let  $R_n$  denote the corresponding right-endpoint sum. In the following exercises, compute the indicated left and right sums for the given functions on the indicated interval.

$$R_4 \text{ for } \frac{1}{x^2 + 1} \text{ on } [-2, 2]$$

### Solution

Since we're using the right-endpoint sum with  $n = 4$  to approximate the integral of  $1/(x^2 + 1)$  from  $-2$  to  $2$ , the sum is taken from 1 to 4 rather than 0 to 3.

$$\begin{aligned} \int_{-2}^2 \frac{1}{x^2 + 1} dx &\approx \sum_{i=1}^4 \frac{1}{x_i^2 + 1} \Delta x = \sum_{i=1}^4 \frac{1}{(-2 + i\Delta x)^2 + 1} \Delta x \\ &= \sum_{i=1}^4 \frac{1}{[4 - 2i\Delta x + i^2(\Delta x)^2] + 1} \Delta x \\ &= \sum_{i=1}^4 \frac{1}{5 - 2i\Delta x + i^2(\Delta x)^2} \Delta x \\ &= \sum_{i=1}^4 \frac{1}{5 - 2i \left[ \frac{2 - (-2)}{4} \right] + i^2 \left[ \frac{2 - (-2)}{4} \right]^2} \left[ \frac{2 - (-2)}{4} \right] \\ &= \sum_{i=1}^4 \frac{1}{5 - 2i(1) + i^2(1)^2} (1) \\ &= \sum_{i=1}^4 \frac{1}{5 - 2i + i^2} \\ &= \frac{1}{5 - 2(1) + (1)^2} + \frac{1}{5 - 2(2) + (2)^2} + \frac{1}{5 - 2(3) + (3)^2} + \frac{1}{5 - 2(4) + (4)^2} \\ &= \frac{1}{4} + \frac{1}{5} + \frac{1}{8} + \frac{1}{13} \\ &= \frac{339}{520} \end{aligned}$$